



Robotics - Homogeneous coordinates and transformations

Simone Ceriani

ceriani@elet.polimi.it

Dipartimento di Elettronica e Informazione
Politecnico di Milano

15 March 2012

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry

⑧ Projective Transformations



Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry

⑧ Projective Transformations



Calendar

1ST PART

WED 14/03 Homogeneous coordinate

THU 29/03 Computer Vision (1)


THU 12/04 Computer Vision (2)

2ND PART

THU 10/05 Localization (1)

THU 07/06 Localization (2)

THU 14/06 Slam



Outline

1 Introduction

2 2D space

3 3D space

4 Rototranslation - 2D

5 Rototranslation - 3D

6 Composition

7 Projective 2D Geometry

8 Projective Transformations



Homogeneous coordinates

INTRODUCTION

- Introduced in 1827 (Möbius)
- Used in *projective geometry*
- Suitable for *points at the infinity*
- Easily code
 - points (2D-3D)
 - lines (2D)
 - conics (2D)
 - planes (3D)
 - quadrics (3D)
 - ...
- Transformation simpler than Cartesian
- ...



A. F. Möbius.


Points in Homogeneous coordinates - 2D space - Definition

HOMOGENEOUS 2D SPACE

- Given a point $\mathbf{p}_e = \begin{bmatrix} X \\ Y \end{bmatrix} \in \mathbb{R}^2$ in Cartesian coordinates
- we can define $\mathbf{p}_h = \begin{bmatrix} x \\ y \\ w \end{bmatrix} \in \mathbb{R}^3$ in homogeneous coordinates
- under the relation $\begin{cases} X &= x/w \\ Y &= y/w \\ w &\neq 0 \end{cases}$
- i.e., there is an arbitrary *scale factor* (w)

Points in Homogeneous coordinates - 2D space - Example

EXAMPLE




- $p_e = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ (euclidean)
- $p_{h_1} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \equiv p_e$
- $p_{h_2} = \begin{bmatrix} 6 \\ 4 \\ 2 \end{bmatrix} \equiv p_e$
- $p_{h_3} = \begin{bmatrix} 1.5 \\ 1 \\ 0.5 \end{bmatrix} \equiv p_e$

- $p_{h_1} \equiv p_{h_2} \equiv p_{h_3}$

Points in Homogeneous coordinates - 2D space - Example

EXAMPLE



- $p_e = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ (euclidean)

- $p_{h_1} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \equiv p_e$

- $p_{h_2} = \begin{bmatrix} 6 \\ 4 \\ 2 \end{bmatrix} \equiv p_e$

- $p_{h_3} = \begin{bmatrix} 1.5 \\ 1 \\ 0.5 \end{bmatrix} \equiv p_e$

- $p_{h_1} \equiv p_{h_2} \equiv p_{h_3}$

NOTE

A Cartesian point can be represented by infinitely many homogeneous coordinates

Points in Homogeneous coordinates - 2D space - Properties

NOTE

A Cartesian point can be represented by infinitely many homogeneous coordinates

PROPERTY

- given $\mathbf{p}_h = \begin{bmatrix} x \\ y \\ w \end{bmatrix}, w \neq 0$
- for $\forall \lambda \neq 0 \quad \hat{\mathbf{p}}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \equiv \mathbf{p}_h$

Points in Homogeneous coordinates - 2D space - Properties

NOTE

A Cartesian point can be represented by infinitely many homogeneous coordinates

PROPERTY

- given $\mathbf{p}_h = \begin{bmatrix} x \\ y \\ w \end{bmatrix}, w \neq 0$
 - for $\forall \lambda \neq 0$ $\hat{\mathbf{p}}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \equiv \mathbf{p}_h$

PROOF

- $\mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$
- for $\forall \lambda \neq 0$ $\hat{\mathbf{p}}_e = \begin{bmatrix} \frac{\lambda x}{\lambda w} \\ \frac{\lambda y}{\lambda w} \end{bmatrix} = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$

Points in Homogeneous coordinates - 2D space - Properties

NOTE

A Cartesian point can be represented by infinitely many homogeneous coordinates

PROPERTY

- given $\mathbf{p}_h = \begin{bmatrix} x \\ y \\ w \end{bmatrix}, w \neq 0$

- for $\forall \lambda \neq 0$ $\hat{\mathbf{p}}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \equiv \mathbf{p}_h$

PROOF

- $\mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$

- for $\forall \lambda \neq 0$ $\hat{\mathbf{p}}_e = \begin{bmatrix} \frac{\lambda x}{\lambda w} \\ \frac{\lambda y}{\lambda w} \end{bmatrix} = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$

NOTES

- $w = 1$: *normalized* homogeneous coordinates
- normalization* : $[x \ y \ w]^T \rightarrow [x/w, y/w, 1]^T, w \neq 0$
- hom* \rightarrow *cart* : $[x \ y \ w]^T \rightarrow [x/w, y/w]^T, w \neq 0$
- cart* \rightarrow *hom* : $[x \ y]^T \rightarrow [x, y, 1]^T$

Points in Homogeneous coordinates - 2D space - Improper points

WHAT'S MORE THAN CARTESIAN?

- All Cartesian points can be expressed in homogeneous coordinates: $\mathbf{p}_e \rightarrow [\mathbf{p}_e, 1]^T$
 - Are homogeneous coordinates more powerful than Cartesian ones? \rightarrow YES

Points in Homogeneous coordinates - 2D space - Improper points

WHAT'S MORE THAN CARTESIAN?

- All Cartesian points can be expressed in homogeneous coordinates: $\mathbf{p}_e \rightarrow [\mathbf{p}_e, 1]^T$
- Are homogeneous coordinates more powerful than Cartesian ones? \rightarrow YES

IMPROPER POINTS

- With $w = 0$ we can express *points at the infinity* $\rightarrow [x/0, y/0]^T$
- $\mathbf{p}_h = [x, y, 0]^T$ codes a *direction*
not directly expressed in Cartesian coordinates

Points in Homogeneous coordinates - 2D space - Improper points

WHAT'S MORE THAN CARTESIAN?

- All Cartesian points can be expressed in homogeneous coordinates: $\mathbf{p}_e \rightarrow [\mathbf{p}_e, 1]^T$
- Are homogeneous coordinates more powerful than Cartesian ones? \rightarrow YES

IMPROPER POINTS


- With $w = 0$ we can express *points at the infinity* $\rightarrow [x/0, y/0]^T$
- $\mathbf{p}_h = [x, y, 0]^T$ codes a *direction*
not directly expressed in Cartesian coordinates

PROPERTY

- $\mathbf{p}_h = [x, y, 0]^T \equiv [\lambda x, \lambda y, 0]^T \forall \lambda \neq 0$

Points in Homogeneous coordinates - 2D space - Directions Example


EXAMPLE



- $\mathbf{d}_x = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$: x-axis
- $\mathbf{d}_y = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$: y-axis
- $\mathbf{d}_{45^\circ} = \begin{bmatrix} \sqrt{2}/2 \\ \sqrt{2}/2 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$: 45° axis

Points in Homogeneous coordinates - 2D space - Directions Example

EXAMPLE



- $d_x = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$: x-axis
- $d_y = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$: y-axis
- $d_{45^\circ} = \begin{bmatrix} \sqrt{2}/2 \\ \sqrt{2}/2 \\ 0 \end{bmatrix} \equiv \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$: 45° axis

NOTE

- A direction can be represented by infinitely many homogeneous directions
- A *unit vector* is the direction with $\|d\| = 1$ (i.e., $\sqrt{x^2 + y^2} = 1$)

Points in Homogeneous coordinates - 2D space - Final Remarks

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

Points in Homogeneous coordinates - 2D space - Final Remarks

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

Points in Homogeneous coordinates - 2D space - Final Remarks

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

IMPROPER POINTS - DIRECTIONS

$$\mathbf{d}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ 0 \end{bmatrix}$$

with $(x \neq 0 \parallel y \neq 0) \ \&\& \ \lambda \neq 0$

Points in Homogeneous coordinates - 2D space - Final Remarks

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

IMPROPER POINTS - DIRECTIONS

$$\mathbf{d}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ 0 \end{bmatrix}$$

with $(x \neq 0 \parallel y \neq 0) \ \&& \ \lambda \neq 0$

INVALID HOMOGENEOUS POINT

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

→ homogeneous 2D space is defined
on $\mathbb{R}^3 - [0, 0, 0]^T$

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry

⑧ Projective Transformations



Points in Homogeneous coordinates - 3D space - Definition

HOMOGENEOUS 3D SPACE

- Given a point $\mathbf{p}_e = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \in \mathbb{R}^3$ in Cartesian coordinates
 - we can define $\mathbf{p}_h = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \in \mathbb{R}^4$ in homogeneous coordinates
 - under the relation $\begin{cases} X &= x/w \\ Y &= y/w \\ Z &= z/w \\ w &\neq 0 \end{cases}$
 - i.e., there is an arbitrary *scale factor* (w)

Points in Homogeneous coordinates - 3D space - Summary

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \\ z/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

Points in Homogeneous coordinates - 3D space - Summary

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \\ z/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

Points in Homogeneous coordinates - 3D space - Summary

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \\ z/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

IMPROPER POINTS - DIRECTIONS

$$\mathbf{d}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ 0 \end{bmatrix} \text{ with}$$

$(x \neq 0 \parallel y \neq 0 \parallel z \neq 0) \And \lambda \neq 0$

Points in Homogeneous coordinates - 3D space - Summary

POINTS

$$\mathbf{p}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ \lambda w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} x/w \\ y/w \\ z/w \end{bmatrix}$$

with $w \neq 0, \lambda \neq 0$

ORIGIN

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ 0 \\ w \end{bmatrix} \rightarrow \mathbf{p}_e = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

with $w \neq 0$

IMPROPER POINTS - DIRECTIONS

$$\mathbf{d}_h = \begin{bmatrix} \lambda x \\ \lambda y \\ \lambda z \\ 0 \end{bmatrix} \text{ with}$$

$(x \neq 0 \parallel y \neq 0 \parallel z \neq 0) \ \&\ \lambda \neq 0$

INVALID HOMOGENEOUS POINT

$$\mathbf{p}_h = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

\rightarrow homogeneous 3D space is defined
on $\mathbb{R}^4 - [0, 0, 0, 0]^T$

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry


⑧ Projective Transformations



Translation - Cartesian 2D


START WITH AN EXAMPLE

- $\mathbf{p}_1^{(0_1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$



Translation - Cartesian 2D


START WITH AN EXAMPLE



- $p_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$
- $O_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \equiv p_1^{(O_1)}$

Translation - Cartesian 2D


START WITH AN EXAMPLE



- $p_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$
- $O_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \equiv p_1^{(O_1)}$
- $p_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \end{bmatrix}$

Translation - Cartesian 2D


START WITH AN EXAMPLE



- $\mathbf{p}_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$
- $O_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \equiv \mathbf{p}_1^{(O_1)}$
- $\mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \end{bmatrix}$
- $\mathbf{p}_2^{(O_1)} = \mathbf{p}_1^{(O_1)} + \mathbf{p}_2^{(O_2)}$
 $= \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix}$

Translation - Homogeneous 2D - 1


SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATE



$$\bullet \quad p_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 & 2 \\ \lambda_1 & 1 \\ \lambda_1 & 1 \end{bmatrix}$$

Translation - Homogeneous 2D - 1


SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATE



$$\bullet \quad \mathbf{p}_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$$

Translation - Homogeneous 2D - 1

SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATES




$$\bullet \quad \mathbf{p}_1^{(o_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$$

$$\bullet \quad O_2 \quad \equiv \mathbf{p}_1^{(O_1)}$$

$$\bullet \quad \mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$$

Translation - Homogeneous 2D - 1

SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATES




$$\bullet \quad \mathbf{p}_1^{(0_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$$

$$\bullet \quad \mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 \cdot 1.5 \\ \lambda_2 \cdot 1.5 \\ \lambda_2 \cdot 1 \end{bmatrix}$$

$$\bullet \quad p_2^{(O_1)} = p_1^{(O_1)} + p_2^{(O_2)}$$

Translation - Homogeneous 2D - 1


SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATES



- $\mathbf{p}_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$
 - $O_2 \equiv \mathbf{p}_1^{(O_1)}$
 - $\mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$
 - ~~$\mathbf{p}_2^{(O_1)} = \mathbf{p}_1^{(O_1)} + \mathbf{p}_2^{(O_2)}$~~ → NO
 - valid only if $\mathbf{p}_1^{(O_1)}_w = \mathbf{p}_2^{(O_2)}_w$
 - e.g., $\begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} + \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \begin{bmatrix} 1.6 \\ 1.3 \\ 1 \end{bmatrix} \not\equiv \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$

Translation - Homogeneous 2D - 1

SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATES




- $\mathbf{p}_1^{(0_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$
 - $O_2 \equiv \mathbf{p}_1^{(0_1)}$
 - $\mathbf{p}_2^{(0_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$
 - $\mathbf{p}_2^{(0_1)} = \mathbf{p}_1^{(0_1)} + \mathbf{p}_2^{(0_2)} \rightarrow \text{NO}$

- valid only if $\frac{\mathbf{p}_1^{(O_1)}}{w} = \frac{\mathbf{p}_2^{(O_2)}}{w}$
 - e.g.,
$$\begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} + \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \begin{bmatrix} 1.6 \\ 1.3 \\ 1 \end{bmatrix} \not\equiv \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$$

- \rightarrow we can normalize points ($w = 1$)
 - e.g.,
$$\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$$

Translation - Homogeneous 2D - 1

SAME EXAMPLE BUT WITH HOMOGENEOUS COORDINATES



- $\mathbf{p}_1^{(0_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$
 - $O_2 \equiv \mathbf{p}_1^{(0_1)}$
 - $\mathbf{p}_2^{(0_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$
 - $\mathbf{p}_2^{(0_1)} = \mathbf{p}_1^{(0_1)} + \mathbf{p}_2^{(0_2)} \rightarrow \text{NO}$

Anything better?

- valid only if $\mathbf{p}_1^{(O_1)}_w = \mathbf{p}_2^{(O_2)}_w$
 - e.g.,
$$\begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} + \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \begin{bmatrix} 1.6 \\ 1.3 \\ 1 \end{bmatrix} \neq \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$$
 - \rightarrow we can normalize points ($w = 1$)
 - e.g.,
$$\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$$

Translation - Homogeneous 2D - 2


TRANSLATION: THE RIGHT WAY WITH HOMOGENEOUS COORDINATES

- $\mathbf{O} = \begin{bmatrix} x_O \\ y_O \\ 1 \end{bmatrix}$: position of the second reference frame (*normalized*)
 - $\mathbf{p}^{(O)} = \begin{bmatrix} x_p \\ y_p \\ w_p \end{bmatrix}$: point w.r.t. the second reference frame (*homogeneous*)

$$\mathbf{p} = \begin{bmatrix} 1 & 0 & x_0 \\ 0 & 1 & y_0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_p \\ y_p \\ w_p \end{bmatrix} = \begin{bmatrix} x_p + x_0 w_p \\ y_p + y_0 w_p \\ w_p \end{bmatrix}$$

Translation - Homogeneous 2D - 3


LET'S TRY ON THE EXAMPLE



- $\mathbf{p}_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$
- $O_2 \equiv \mathbf{p}_1^{(O_1)}$ normalized
- $\mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$

Translation - Homogeneous 2D - 3

LET'S TRY ON THE EXAMPLE




- $\mathbf{p}_1^{(O_1)} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_1 2 \\ \lambda_1 1 \\ \lambda_1 1 \end{bmatrix}$
- $O_2 \equiv \mathbf{p}_1^{(O_1)}$ normalized
- $\mathbf{p}_2^{(O_2)} = \begin{bmatrix} 1.5 \\ 1.5 \\ 1 \end{bmatrix} \equiv \begin{bmatrix} 6 \\ 6 \\ 4 \end{bmatrix} \equiv \dots \equiv \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix}$

$$\begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \lambda_2 1.5 \\ \lambda_2 1.5 \\ \lambda_2 1 \end{bmatrix} = \begin{bmatrix} \lambda_2 1.5 + \lambda_2 2 \\ \lambda_2 1.5 + \lambda_2 1 \\ \lambda_2 \end{bmatrix} \equiv \begin{bmatrix} 3.5 \\ 2.5 \\ 1 \end{bmatrix}$$

Rotation - Cartesian 2D


DERIVATION FROM EXAMPLE

- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$



Rotation - Cartesian 2D


DERIVATION FROM EXAMPLE



- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$

Rotation - Cartesian 2D


DERIVATION FROM EXAMPLE



- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$
- $\mathbf{p}^{(O_2)} = \begin{bmatrix} \mathbf{p}_x \\ \mathbf{p}_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$

Rotation - Cartesian 2D


DERIVATION FROM EXAMPLE



- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$
- $p^{(O_2)} = \begin{bmatrix} p_x \\ p_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$

Rotation - Cartesian 2D


DERIVATION FROM EXAMPLE



- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$
- $\mathbf{p}^{(O_2)} = \begin{bmatrix} \mathbf{p}_x \\ \mathbf{p}_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$
- $\cos(\theta + 90^\circ) = -\sin(\theta)$
- $\sin(\theta + 90^\circ) = \cos(\theta)$

Rotation - Cartesian 2D

DERIVATION FROM EXAMPLE



- $O_1 \equiv O_2$
- rotated of $\theta = 30^\circ$
- $\mathbf{p}^{(O_2)} = \begin{bmatrix} \mathbf{p}_x \\ \mathbf{p}_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$
- $\cos(\theta + 90^\circ) = -\sin(\theta)$
- $\sin(\theta + 90^\circ) = \cos(\theta)$

$$\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta)\mathbf{p}_x - \sin(\theta)\mathbf{p}_y \\ \sin(\theta)\mathbf{p}_x + \cos(\theta)\mathbf{p}_y \end{bmatrix} = \begin{bmatrix} \cos(30^\circ)2 - \sin(30^\circ)2 \\ \sin(20^\circ)2 + \cos(30^\circ)2 \end{bmatrix} = \begin{bmatrix} 0.73 \\ 2.73 \end{bmatrix}$$

Rotation - Homogeneous 2D

- From $\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta)\mathbf{p}_x - \sin(\theta)\mathbf{p}_y \\ \sin(\theta)\mathbf{p}_x + \cos(\theta)\mathbf{p}_y \end{bmatrix}$

Rotation - Homogeneous 2D


- From $\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta)\mathbf{p}_x - \sin(\theta)\mathbf{p}_y \\ \sin(\theta)\mathbf{p}_x + \cos(\theta)\mathbf{p}_y \end{bmatrix}$
- Rewrite with matrices $\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \mathbf{p}^{(O_2)}$

Rotation - Homogeneous 2D

- From $\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta)\mathbf{p}_x - \sin(\theta)\mathbf{p}_y \\ \sin(\theta)\mathbf{p}_x + \cos(\theta)\mathbf{p}_y \end{bmatrix}$
- Rewrite with matrices $\mathbf{p}^{(O_1)} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \mathbf{p}^{(O_2)}$
- Pass to homogeneous $\mathbf{p}_h^{(O_1)} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{p}_h^{(O_2)}$
 where $\mathbf{p}_h^{(O_2)} = \begin{bmatrix} \lambda\mathbf{p}_x \\ \lambda\mathbf{p}_y \\ \lambda \end{bmatrix}$

Rototranslation - Homogeneous 2D

PUTTING THINGS TOGETHER



- O_2 translated of \mathbf{t} w.r.t. O_1
- O_2 rotated of θ w.r.t. O_1

TWO STEPS:

$$\mathbf{p}_h^{(O'_2)} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{p}_h^{(O_2)}$$

$$\mathbf{p}_h^{(O_1)} = \begin{bmatrix} 1 & 0 & \mathbf{t}_x \\ 0 & 1 & \mathbf{t}_y \\ 0 & 0 & 1 \end{bmatrix} \mathbf{p}_h^{(O'_2)}$$

ONE STEP:

$$\mathbf{p}_h^{(O^1)} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & \mathbf{t}_x \\ \sin(\theta) & \cos(\theta) & \mathbf{t}_y \\ 0 & 0 & 1 \end{bmatrix} \mathbf{p}_h^{(O_2)}$$

Rototranslation - Homogeneous 2D - Some notes

$$\text{Consider } \begin{bmatrix} \cos(\theta) & -\sin(\theta) & t_x \\ \sin(\theta) & \cos(\theta) & t_y \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} n_x & m_x & t_x \\ n_y & m_y & t_y \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix} = T$$

- $\mathbf{n} = \begin{bmatrix} n_x \\ n_y \\ 0 \end{bmatrix}$ is the *direction vector* (improper point) of the x axis
 - $\mathbf{m} = \begin{bmatrix} m_x \\ m_y \\ 0 \end{bmatrix}$ is the *direction vector* (improper point) of the x axis
 - $\|\mathbf{n}\| = \|\mathbf{m}\| = 1$ they are unit vectors
 - $\|[\mathbf{n}_x, \mathbf{m}_x]\| = \|[\mathbf{n}_y, \mathbf{m}_y]\| = 1$ are unit vectors too
 - \mathbf{R} is an *orthogonal matrix* $\rightarrow \mathbf{R}^{-1} = \mathbf{R}^T$
 - \mathbf{T} is homogeneous too! i.e., $\mathbf{T} \equiv \lambda \mathbf{T} \quad \mathbf{T} \mathbf{p}_2 \equiv \lambda \mathbf{T} \mathbf{p}_2$

Rototranslation - Homogeneous 2D - Get parameters

Consider $\mathbf{T} = \begin{bmatrix} T_{11} & T_{12} & T_{13} \\ T_{21} & T_{22} & T_{23} \\ 0 & 0 & 1 \end{bmatrix}$

- remember $\begin{bmatrix} \cos(\theta) & -\sin(\theta) & t_x \\ \sin(\theta) & \cos(\theta) & t_y \\ 0 & 0 & 1 \end{bmatrix}$

- $\mathbf{t} = \begin{bmatrix} T_{13} \\ T_{23} \end{bmatrix}$

- $\theta = \text{atan2}(T_{21}, T_{11})$


Rototranslation - Homogeneous 2D - Get parameters

Consider $\mathbf{T} = \begin{bmatrix} T_{11} & T_{12} & T_{13} \\ T_{21} & T_{22} & T_{23} \\ 0 & 0 & 1 \end{bmatrix}$

- remember $\begin{bmatrix} \cos(\theta) & -\sin(\theta) & t_x \\ \sin(\theta) & \cos(\theta) & t_y \\ 0 & 0 & 1 \end{bmatrix}$

- $\mathbf{t} = \begin{bmatrix} T_{13} \\ T_{23} \end{bmatrix}$

- $\theta = \text{atan2}(T_{21}, T_{11})$



- $\arctan(y/x) \Rightarrow [0, \pi]$
- $\text{atan2}(y, x) \rightarrow [-\pi, \pi]$


Rototranslation - Homogeneous 2D - Get parameters

$$\text{Consider } \mathbf{T} = \begin{bmatrix} T_{11} & T_{12} & T_{13} \\ T_{21} & T_{22} & T_{23} \\ 0 & 0 & 1 \end{bmatrix}$$

- remember $\begin{bmatrix} \cos(\theta) & -\sin(\theta) & t_x \\ \sin(\theta) & \cos(\theta) & t_y \\ 0 & 0 & 1 \end{bmatrix}$

- $t = \begin{bmatrix} T_{13} \\ T_{23} \end{bmatrix}$

- $\theta = \text{atan2}(T_{21}, T_{11})$



- $\arctan(y/x) \rightarrow [0, \pi]$
 - $\text{atan2}(y, x) \rightarrow [-\pi, \pi]$

$$\text{atan2}(y, x) = \begin{cases} \arctan(y/x) & x > 0 \\ \arctan(y/x) + \pi & y \geq 0, x < 0 \\ \arctan(y/x) - \pi & y < 0, x < 0 \\ +\frac{\pi}{2} & y > 0, x = 0 \\ -\frac{\pi}{2} & y < 0, x = 0 \\ \text{undefined} & x = 0, y = 0 \end{cases}$$

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition


⑦ Projective 2D Geometry

⑧ Projective Transformations



Rotations in 3D - 1

ROTATE AROUND X




ρ , roll ("rollio")
around x-axis

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\rho) & -\sin(\rho) \\ 0 & \sin(\rho) & \cos(\rho) \end{bmatrix}$$


Rotations in 3D - 1

ROTATE AROUND X



ρ , *roll* ("rollio")
around x -axis

ROTATE AROUND Y




θ , *pitch* ("beccheggio")
around y -axis

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\rho) & -\sin(\rho) \\ 0 & \sin(\rho) & \cos(\rho) \end{bmatrix} \quad \mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$


Rotations in 3D - 1

ROTATE AROUND X




ρ , *roll* ("rollio")
around x -axis

ROTATE AROUND Y



θ , *pitch* ("beccheggio")
around y -axis

ROTATE AROUND Z



ϕ , *yaw* ("imbardata")
around z -axis

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\rho) & -\sin(\rho) \\ 0 & \sin(\rho) & \cos(\rho) \end{bmatrix}$$

$$\mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

$$\mathbf{R}_z = \begin{bmatrix} \cos(\phi) & -\sin(\phi) & 0 \\ \sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Rotations in 3D - 2

To CREATE A 3D ROTATION

- Compose 3 planar rotation
- 24 possible conventions
 - non-commutative matrix products

COMMON CONVENTION

- Rotate around x (roll)
- Rotate around y (pitch)
- Rotate around z (yaw)

Rotations in 3D - 2

TO CREATE A 3D ROTATION

- Compose 3 planar rotation
- 24 possible conventions
non-commutative matrix products

COMMON CONVENTION

- Rotate around x (roll)
- Rotate around y (pitch)
- Rotate around z (yaw)

DERIVE A COMPLETE ROTATION MATRIX

- Let \mathbf{p}^{Oxyz} in the rotated system
- $\mathbf{p}^{Oxyz} = \mathbf{R}_x \mathbf{p}^{Oxyz}$
- $\mathbf{p}^{Oz} = \mathbf{R}_y \mathbf{p}^{Oxyz}$
- $\mathbf{p}^O = \mathbf{R}_z \mathbf{p}^{Oxyz}$ in the original system

Rotations in 3D - 2

TO CREATE A 3D ROTATION

- Compose 3 planar rotation
- 24 possible conventions
non-commutative matrix products

COMMON CONVENTION

- Rotate around x (roll)
- Rotate around y (pitch)
- Rotate around z (yaw)

DERIVE A COMPLETE ROTATION MATRIX

- Let $\mathbf{p}^{O^R_{xyz}}$ in the rotated system
- $\mathbf{p}^{O^R_{yz}} = \mathbf{R}_x \mathbf{p}^{O^R_{xyz}}$
- $\mathbf{p}^{O^R_z} = \mathbf{R}_y \mathbf{p}^{O^R_{yz}}$
- $\mathbf{p}^O = \mathbf{R}_z \mathbf{p}^{O^R_z}$ in the original system
- $\mathbf{R}_{xyz} = \mathbf{R}_z \mathbf{R}_y \mathbf{R}_x$

Rotations in 3D - 2

TO CREATE A 3D ROTATION

- Compose 3 planar rotation
- 24 possible conventions
non-commutative matrix products

COMMON CONVENTION


- Rotate around x (roll)
- Rotate around y (pitch)
- Rotate around z (yaw)

DERIVE A COMPLETE ROTATION MATRIX

- Let $\mathbf{p}^{O^R_{xyz}}$ in the rotated system
- $\mathbf{p}^{O^R_{yz}} = \mathbf{R}_x \mathbf{p}^{O^R_{xyz}}$
- $\mathbf{p}^{O^R_z} = \mathbf{R}_y \mathbf{p}^{O^R_{yz}}$
- $\mathbf{p}^O = \mathbf{R}_z \mathbf{p}^{O^R_z}$ in the original system
- $\mathbf{R}_{xyz} = \mathbf{R}_z \mathbf{R}_y \mathbf{R}_x$

$$\mathbf{R}_{xyz} = \begin{bmatrix} \cos(\phi) \cos(\theta) & \cos(\phi) \sin(\theta) \sin(\rho) - \sin(\phi) \cos(\rho) & \cos(\phi) \sin(\theta) \cos(\rho) + \sin(\phi) \sin(\rho) \\ \sin(\phi) \cos(\theta) & \sin(\phi) \sin(\theta) \sin(\rho) + \cos(\phi) \cos(\rho) & \sin(\phi) \sin(\theta) \cos(\rho) - \cos(\phi) \sin(\rho) \\ -\sin(\theta) & \cos(\theta) \sin(\rho) & \cos(\theta) \cos(\rho) \end{bmatrix}$$


Rototranslations - Homogeneous 3D



COMPLETE TRANSFORMATION

- $\mathbf{p}^{(O_2)}$ w.r.t. O_2 reference
- Let consider O'_2 rotated as O_1 , but translated by \mathbf{t}
- \mathbf{R} rotation of O_2 wrt O'_2


Rototranslations - Homogeneous 3D



COMPLETE TRANSFORMATION

- $\mathbf{p}^{(O_2)}$ w.r.t. O_2 reference
- Let consider O'_2 rotated as O_1 , but translated by \mathbf{t}
- \mathbf{R} rotation of O_2 wrt O'_2
- $\mathbf{p}^{(O'_2)} = \mathbf{R} \mathbf{p}^{(O_2)}$


Rototranslations - Homogeneous 3D



COMPLETE TRANSFORMATION

- $\mathbf{p}^{(O_2)}$ w.r.t. O_2 reference
- Let consider O'_2 rotated as O_1 , but translated by \mathbf{t}
- \mathbf{R} rotation of O_2 wrt O'_2
- $\mathbf{p}^{(O'_2)} = \mathbf{R}\mathbf{p}^{(O_2)}$
- $\mathbf{p}^{(O_1)} = \mathbf{t} + \mathbf{p}^{(O'_2)}$

Rototranslations - Homogeneous 3D



COMPLETE TRANSFORMATION

- $\mathbf{p}^{(O_2)}$ w.r.t. O_2 reference
- Let consider O'_2 rotated as O_1 , but translated by \mathbf{t}
- \mathbf{R} rotation of O_2 wrt O'_2
- $\mathbf{p}^{(O'_2)} = \mathbf{R}\mathbf{p}^{(O_2)}$
- $\mathbf{p}^{(O_1)} = \mathbf{t} + \mathbf{p}^{(O'_2)}$

IN HOMOGENEOUS COORDINATES

$$\mathbf{p}_h^{(O_1)} = \begin{bmatrix} \mathbf{I} & \mathbf{t} \\ \mathbf{0} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{R} & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} \mathbf{p}_h^{(O_2)} = \begin{bmatrix} \mathbf{R} & \mathbf{t} \\ \mathbf{0} & 1 \end{bmatrix} \mathbf{p}_h^{(O_2)}$$

where $\mathbf{p}_h^{(O_2)}$ is $\mathbf{p}^{(O_2)}$ in homogeneous coordinates

and $\mathbf{p}_h^{(O_1)}$ is $\mathbf{p}^{(O_1)}$ in homogeneous coordinates

Rototranslation - Homogeneous 3D - Some notes

Consider

$$\begin{bmatrix} \mathbf{n}_x & \mathbf{m}_x & \mathbf{a}_x & t_x \\ \mathbf{n}_y & \mathbf{m}_y & \mathbf{a}_y & t_y \\ \mathbf{n}_z & \mathbf{m}_z & \mathbf{a}_z & t_z \\ 0 & 0 & 1 & \end{bmatrix} = \begin{bmatrix} \mathbf{R} & \mathbf{t} \\ \mathbf{0} & 1 \end{bmatrix} = \mathbf{T}$$

- $\mathbf{n} = \begin{bmatrix} \mathbf{n}_x \\ \mathbf{n}_y \\ \mathbf{n}_z \\ 0 \end{bmatrix}$, $\mathbf{m} = \begin{bmatrix} \mathbf{m}_x \\ \mathbf{m}_y \\ \mathbf{m}_z \\ 0 \end{bmatrix}$, $\mathbf{a} = \begin{bmatrix} \mathbf{a}_x \\ \mathbf{a}_y \\ \mathbf{a}_z \\ 0 \end{bmatrix}$ are the *direction vectors* of the x,y,z , axes
- $\|\mathbf{n}\| = \|\mathbf{m}\| = \|\mathbf{a}\| = 1$ unit vectors
- $\|[\mathbf{n}_x, \mathbf{m}_x, \mathbf{a}_x]\| = \|[\mathbf{n}_y, \mathbf{m}_y, \mathbf{a}_y]\| = \|[\mathbf{n}_z, \mathbf{m}_z, \mathbf{a}_z]\| = 1$ are unit vectors too
- \mathbf{R} is an *orthogonal matrix* $\rightarrow \mathbf{R}^{-1} = \mathbf{R}^T$
- \mathbf{T} is homogeneous too! i.e., $\mathbf{T} \equiv \lambda \mathbf{T} \quad \mathbf{T} \mathbf{p}_2 \equiv \lambda \mathbf{T} \mathbf{p}_2$

Rototranslation - Homogeneous 3D - Get parameters

$$\text{Consider } \mathbf{T} = \begin{bmatrix} T_{11} & T_{12} & T_{13} & T_{14} \\ T_{21} & T_{22} & T_{23} & T_{24} \\ T_{31} & T_{32} & T_{33} & T_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- remember $\begin{bmatrix} \cos(\phi) \cos(\theta) & \cos(\phi) \sin(\theta) \sin(\rho) - \sin(\phi) \cos(\rho) & \cos(\phi) \sin(\theta) \cos(\rho) + \sin(\phi) \sin(\rho) \\ \sin(\phi) \cos(\theta) & \sin(\phi) \sin(\theta) \sin(\rho) + \cos(\phi) \cos(\rho) & \sin(\phi) \sin(\theta) \cos(\rho) - \cos(\phi) \sin(\rho) \\ -\sin(\theta) & \cos(\theta) \sin(\rho) & \cos(\theta) \cos(\rho) \end{bmatrix}$

- $\mathbf{t} = \begin{bmatrix} T_{14} \\ T_{24} \\ T_{34} \end{bmatrix}$

- $\phi = \text{atan2}(T_{21}, T_{11})$

- $\theta = \text{atan2}(-T_{31}, \sqrt{T_{32}^2 + T_{33}^2})$

- $\rho = \text{atan2}(T_{32}, T_{33})$

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition ARTIFICIAL INTELLIGENCE and ROBOTICS LABORATORY


⑦ Projective 2D Geometry

⑧ Projective Transformations



Transformations - Why?


THINK ABOUT...



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .

Transformations - Why?


THINK ABOUT...



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .
- The robot perceives the *red point*, it knows the point $p^{(R)}$ in robot reference frame.

Transformations - Why?


THINK ABOUT...



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .
- The robot perceives the *red point*, it knows the point $p^{(R)}$ in robot reference frame.
- $p^{(W)} = T_{WR}^{(W)} p^{(R)}$ is the point in world coordinates.

Transformations - Why?

THINK ABOUT...




- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .
- The robot perceives the *red point*, it knows the point $p^{(R)}$ in robot reference frame.
- $p^{(W)} = T_{WR}^{(W)} p^{(R)}$ is the point in world coordinates.

$$T_{WR}^{(W)} = \begin{bmatrix} R_{WR}^{(W)} & t_{WR}^{(W)} \\ 0 & 1 \end{bmatrix} \text{ is the transformation matrix}$$

- map points in R reference frame in W frame
- $t_{WR}^{(W)}$ is the position of R w.r.t. W
- $R_{WR}^{(W)}$ is the rotation applied to a reference frame rotated as W with origin on O

Transformations - Inversion


INVERSE TRANSFORMATION



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .

Transformations - Inversion

INVERSE TRANSFORMATION



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .


POSITION OF THE WORLD W.R.T. THE ROBOT

$$T_{RW}^{(R)} = \left(T_{WR}^{(W)}\right)^{-1} = \begin{bmatrix} {R_{WR}^{(W)}}^T & -{R_{WR}^{(W)}}^T t_{WR}^{(W)} \\ \mathbf{0} & 1 \end{bmatrix}$$

is the transformation matrix

Transformations - Inversion

INVERSE TRANSFORMATION




- W is the *world* reference frame.
- R is the *robot* reference frame.
- $\mathbf{T}_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .
- You know the *red point* $(\mathbf{p}^{(W)})$ in world coordinates,

POSITION OF THE WORLD W.R.T. THE ROBOT

$$\mathbf{T}_{RW}^{(R)} = (\mathbf{T}_{WR}^{(W)})^{-1} = \begin{bmatrix} \mathbf{R}_{WR}^{(W)^T} & -\mathbf{R}_{WR}^{(W)^T} \mathbf{t}_{WR}^{(W)} \\ \mathbf{0} & 1 \end{bmatrix} \text{ is the transformation matrix}$$

Transformations - Inversion

INVERSE TRANSFORMATION



- W is the *world* reference frame.
- R is the *robot* reference frame.
- $T_{WR}^{(W)}$ is the transformation that codes position and orientation of the robot w.r.t. W .
- You know the *red point* $(p^{(W)})$ in world coordinates,


POSITION OF THE WORLD W.R.T. THE ROBOT

$$T_{RW}^{(R)} = (T_{WR}^{(W)})^{-1} = \begin{bmatrix} R_{WR}^{(W)\top} & -R_{WR}^{(W)\top} t_{WR}^{(W)} \\ \mathbf{0} & 1 \end{bmatrix} \text{ is the transformation matrix}$$

$p^{(R)} = T_{RW}^{(R)} p^{(W)}$ is the point in robot coordinates.

Transformations - Composition


COMPOSITION OF TRANSFORMATIONS



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1

Transformations - Composition


COMPOSITION OF TRANSFORMATIONS



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1
- $T_{02}^{(0)} = T_{01}^{(0)} T_{12}^{(1)}$: pose of 2 w.r.t. 0

Transformations - Composition


COMPOSITION OF TRANSFORMATIONS



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1
- $T_{02}^{(0)} = T_{01}^{(0)} T_{12}^{(1)}$: pose of 2 w.r.t. 0
- $T_{23}^{(2)}$: pose of 3 w.r.t. 2

Transformations - Composition


COMPOSITION OF TRANSFORMATIONS



- $\mathbf{T}_{01}^{(0)}$: pose of 1 w.r.t. 0
- $\mathbf{T}_{12}^{(1)}$: pose of 2 w.r.t. 1
- $\mathbf{T}_{02}^{(0)} = \mathbf{T}_{01}^{(0)} \mathbf{T}_{12}^{(1)}$: pose of 2 w.r.t. 0
- $\mathbf{T}_{23}^{(2)}$: pose of 3 w.r.t. 2
- $\mathbf{T}_{03}^{(0)} = \mathbf{T}_{01}^{(0)} \mathbf{T}_{12}^{(1)} \mathbf{T}_{23}^{(2)} = \mathbf{T}_{02}^{(0)} \mathbf{T}_{23}^{(2)}$: pose of 3 w.r.t. 0

Transformations - Composition

COMPOSITION OF TRANSFORMATIONS



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1
- $T_{02}^{(0)} = T_{01}^{(0)} T_{12}^{(1)}$: pose of 2 w.r.t. 0
- $T_{23}^{(2)}$: pose of 3 w.r.t. 2
- $T_{03}^{(0)} = T_{01}^{(0)} T_{12}^{(1)} T_{23}^{(2)} = T_{02}^{(0)} T_{23}^{(2)}$: pose of 3 w.r.t. 0


GRAPHICAL METHOD

- Post-multiplication following arrows verse
- Pre-multiplication coming back in arrows verse

NOTE: Not unique convention about arrow direction!!

Transformations - Composition Example


EXAMPLE



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1
- $T_{03}^{(0)}$: pose of 3 w.r.t. 0
- $p^{(3)}$: position of p w.r.t 3

Transformations - Composition Example


EXAMPLE



- $T_{01}^{(0)}$: pose of 1 w.r.t. 0
- $T_{12}^{(1)}$: pose of 2 w.r.t. 1
- $T_{03}^{(0)}$: pose of 3 w.r.t. 0
- $p^{(3)}$: position of p w.r.t 3
- $p^{(2)}$: position of p w.r.t 2?

Transformations - Composition Example

EXAMPLE




- $\mathbf{T}_{01}^{(0)}$: pose of 1 w.r.t. 0
- $\mathbf{T}_{12}^{(1)}$: pose of 2 w.r.t. 1
- $\mathbf{T}_{03}^{(0)}$: pose of 3 w.r.t. 0
- $\mathbf{p}^{(3)}$: position of \mathbf{p} w.r.t 3
- $\mathbf{p}^{(2)}$: position of \mathbf{p} w.r.t 2?

SOLUTION

- $\mathbf{T}_{23}^{(2)} = (\mathbf{T}_{12}^{(1)})^{-1} (\mathbf{T}_{01}^{(0)})^{-1} \mathbf{T}_{03}^{(0)} = \mathbf{T}_{21}^{(2)} \mathbf{T}_{10}^{(1)} \mathbf{T}_{03}^{(0)}$
- Note: $(\mathbf{T}_{12}^{(1)})^{-1} (\mathbf{T}_{01}^{(0)})^{-1} = (\mathbf{T}_{01}^{(0)} \mathbf{T}_{12}^{(1)})^{-1} = (\mathbf{T}_{02}^{(0)})^{-1}$
- $\mathbf{p}^{(2)} = \mathbf{T}_{23}^{(2)} \mathbf{p}^{(3)}$

Transformations - Composition - Practical Case


CHANGE REFERENCE SYSTEM OF MOTION



- $\mathbf{T}_{R_0 R_1}^{(R0)}$: pose of robot R at time $t = 1$
w.r.t. robot at time $t = 0$
i.e., the relative motion of the robot
 - $\mathbf{T}_S^{(R)}$: pose of a sensor S w.r.t. the robot R
Note: fixed in time

Transformations - Composition - Practical Case


CHANGE REFERENCE SYSTEM OF MOTION



- $\mathbf{T}_{R_0 R_1}^{(R)}$: pose of robot R at time $t = 1$
w.r.t. robot at time $t = 0$
i.e., the relative motion of the robot
 - $\mathbf{T}_S^{(R)}$: pose of a sensor S w.r.t. the robot R
Note: fixed in time
 - $\mathbf{T}_{S_0 S_1}^{(S_0)}$: pose of sensor S at time $t = 1$
w.r.t. sensor at time $t = 0$?
i.e., the relative motion of the sensor

Transformations - Composition - Practical Case

CHANGE REFERENCE SYSTEM OF MOTION



- $\mathbf{T}_{R_0 R_1}^{(R0)}$: pose of robot R at time $t = 1$
w.r.t. robot at time $t = 0$
i.e., the relative motion of the robot
 - $\mathbf{T}_S^{(R)}$: pose of a sensor S w.r.t. the robot R
Note: fixed in time
 - $\mathbf{T}_{S_0 S_1}^{(S0)}$: pose of sensor S at time $t = 1$
w.r.t. sensor at time $t = 0$?
i.e., the relative motion of the sensor

SOLUTION

$$\bullet \quad \mathbf{T}_{S_0 S_1}^{(S_0)} = \left(\mathbf{T}_{RS}^{(R)} \right)^{-1} \mathbf{T}_{R_0 R_1}^{(R_0)} \mathbf{T}_{RS}^{(R)}$$

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry

⑧ Projective Transformations



Lines in homogeneous coordinates

LINES DEFINITION

- Slope-intercept form: $y = mx + b$

vertical lines $m \equiv \infty$

- Linear equation: $a x + b y + c = 0$,

$$(a, b) \in \mathbb{R}^2 - \{0, 0\}$$

Lines in homogeneous coordinates

HOMOGENEOUS COORDINATES

LINES DEFINITION

- Slope-intercept form: $y = mx + q$

vertical lines $m = \infty$

- Linear equation: $ax + by + c = 0$,


$$(a, b) \in \mathbb{R}^2 - \{0, 0\}$$

- Line: $\mathbf{l} = \begin{bmatrix} a, b, c \end{bmatrix}^T$
- Point: $\mathbf{p} = \begin{bmatrix} x, y, 1 \end{bmatrix}^T$
- \mathbf{p} lies on $\mathbf{l} \Leftrightarrow \mathbf{l}^T \mathbf{p} = \mathbf{p}^T \mathbf{l} = 0$
- Homogeneous property:
 - $\mathbf{l} \equiv \lambda_1 \mathbf{l}$
 - $\mathbf{p} \equiv \lambda_2 \mathbf{p}$

Point from Lines & Lines from Points

INTERSECTION OF LINES

- $\mathbf{p} = \mathbf{l}_1 \cap \mathbf{l}_2 = \mathbf{l}_1 \times \mathbf{l}_2$




- $\mathbf{l}_1 = [1, 0, -1]^T \rightarrow \mathbf{x} = 1$
- $\mathbf{l}_2 = [0, 1, -1]^T \rightarrow \mathbf{y} = 1$
- $\mathbf{p} = [1, 1, 1]^T$

Point from Lines & Lines from Points

INTERSECTION OF LINES


- $\mathbf{p} = \mathbf{l}_1 \cap \mathbf{l}_2 = \mathbf{l}_1 \times \mathbf{l}_2$



- $\mathbf{l}_1 = [1, 0, -1]^T \rightarrow x = 1$
- $\mathbf{l}_2 = [0, 1, -1]^T \rightarrow y = 1$
- $\mathbf{p} = [1, 1, 1]^T$

LINE JOINING TWO POINTS

- $\mathbf{l} = \mathbf{p}_1 \times \mathbf{p}_2$




- $\mathbf{p}_1 = [1, 1, 1]^T$
- $\mathbf{p}_2 = [2, 2, 1]^T$
- $\mathbf{l} = [-1, 1, 0]^T \rightarrow y = x$

Point from Lines & Lines from Points

INTERSECTION OF LINES


- $\mathbf{p} = \mathbf{l}_1 \cap \mathbf{l}_2 = \mathbf{l}_1 \times \mathbf{l}_2$



- $\mathbf{l}_1 = [1, 0, -1]^T \rightarrow x = 1$
- $\mathbf{l}_2 = [0, 1, -1]^T \rightarrow y = 1$
- $\mathbf{p} = [1, 1, 1]^T$

LINE JOINING TWO POINTS

- $\mathbf{l} = \mathbf{p}_1 \times \mathbf{p}_2$



- $\mathbf{p}_1 = [1, 1, 1]^T$
- $\mathbf{p}_2 = [2, 2, 1]^T$
- $\mathbf{l} = [-1, 1, 0]^T \rightarrow y = x$


CROSS PRODUCT REMINDER

$$\mathbf{a} = \begin{bmatrix} a_x \\ a_y \\ a_z \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} b_x \\ b_y \\ b_z \end{bmatrix} \quad \mathbf{a} \times \mathbf{b} = \det \begin{bmatrix} \vec{\mathbf{u}}_x & \vec{\mathbf{u}}_y & \vec{\mathbf{u}}_z \\ \mathbf{a}_x & \mathbf{a}_y & \mathbf{a}_z \\ \mathbf{b}_x & \mathbf{b}_y & \mathbf{b}_z \end{bmatrix} = \begin{bmatrix} \mathbf{a}_y \mathbf{b}_z - \mathbf{a}_z \mathbf{b}_y \\ \mathbf{a}_z \mathbf{b}_x - \mathbf{a}_x \mathbf{b}_z \\ \mathbf{a}_x \mathbf{b}_y - \mathbf{a}_y \mathbf{b}_x \end{bmatrix}$$

Ideal points and \mathbf{l}_∞

INTERSECTION OF PARALLEL LINES

- $\mathbf{p} = \mathbf{l}_1 \cap \mathbf{l}_2 = \mathbf{l}_1 \times \mathbf{l}_2$




- $\mathbf{l}_1 = [1, 0, -1]^T \rightarrow x = 1$
- $\mathbf{l}_2 = [1, 0, -2]^T \rightarrow x = 2$
- $\mathbf{p} = [0, 1, 0]^T \rightarrow$ *improper point*

direction of y axis

Ideal points and I_∞

INTERSECTION OF PARALLEL LINES

- $$\bullet \text{ } P \equiv l_1 \cup l_2 \equiv l_1 \times l_2$$



- $\mathbf{l}_1 = [1, 0, -1]^T \rightarrow x = 1$
 - $\mathbf{l}_2 = [1, 0, -2]^T \rightarrow x = 2$
 - $\mathbf{p} = [0, 1, 0]^T \rightarrow \text{improper point}$
direction of y axis

LINE THAT JOIN IMPROPER POINTS (I_∞)

- $\mathbf{p}_1 = [x_1, y_1, 0]^T$
 - $\mathbf{p}_2 = [x_2, y_2, 0]^T$
 - $\mathbf{p}_1 \times \mathbf{p}_1 \equiv [0, 0, 1]^T$
 - $\mathbf{I}_{\infty} = [0, 0, 1]^T$:

i.e., $\mathbf{I}_{\infty}^T [x, y, 0]^T = 0$

Duality principle

Duality

$$\mathbf{p} \longleftrightarrow \mathbf{l}$$

$$\mathbf{p}^T \mathbf{l} = 0 \longleftrightarrow \mathbf{l}^T \mathbf{p} = 0$$

$$\mathbf{p} = \mathbf{l}_1 \times \mathbf{l}_2 \longleftrightarrow \mathbf{l} = \mathbf{p}_1 \times \mathbf{p}_2$$

To any theorem in 2D projective geometry there correspond a *dual theorem*,
 derived by interchanging the role of *points* and *lines*


Conics

DEFINITION

- $2^n d$ degree equations
- planar curve
- Equation: $ax^2 + bxy + cy^2 + dx + ey + f = 0$
- Homogeneous: $ax^2 + bxy + cy^2 + dxw + eyw + fw^2 = 0$
- Matrix form:
 - $\mathbf{x} = [x, y, w]^T$
 - $\mathbf{C} = \begin{bmatrix} a & b/2 & d/2 \\ b/2 & c & e/2 \\ d/2 & e/2 & f \end{bmatrix}$
 - $\mathbf{x}^T \mathbf{C} \mathbf{x} = 0$

→ \mathbf{C} is homogeneous too, i.e., 6 parameters, 5 D.O.F.


Conics - Summary




PROPER CONICS:

$$\text{rank}(\mathbf{C}) = 3$$


Circle



Ellipse




Hyperbola



Parabola


DEGENERATE CONICS:

$$\text{rank}(\mathbf{C}) < 3$$



$$\mathbf{C} = \mathbf{I}\mathbf{m}^T + \mathbf{m}\mathbf{l}^T$$

2-lines,
 $\text{rank}(\mathbf{C}) = 2$



$$\mathbf{C} = \mathbf{I}\mathbf{l}^T$$

repeated line,
 $\text{rank}(\mathbf{C}) = 1$

Conics Parameters Estimation

PARAMETERS ESTIMATION

- Given a point x_i, y_i , it satisfies

$$ax_i^2 + bx_iy_i + cy_i^2 + dx_i + ey_i + f = 0$$

- Rewrite: $\begin{bmatrix} x_i^2, x_iy_i, y_i^2, x_i, y_i, 1 \end{bmatrix} \begin{bmatrix} a, b, c, d, e, f \end{bmatrix}^T = 0$
- Stacking constraints on ≥ 5 points:

$$\begin{bmatrix} x_1^2 & x_1y_1 & y_1^2 & x_1 & y_1 & 1 \\ x_2^2 & x_2y_2 & y_2^2 & x_2 & y_2 & 1 \\ x_3^2 & x_3y_3 & y_3^2 & x_3 & y_3 & 1 \\ x_4^2 & x_4y_4 & y_4^2 & x_4 & y_4 & 1 \\ x_5^2 & x_5y_5 & y_5^2 & x_5 & y_5 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \\ e \\ f \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Solve the linear system

Details

“Multiple View Geometry in computer vision” - Hartley, Zisserman, Chapter 2.

Outline

① Introduction

② 2D space

③ 3D space

④ Rototranslation - 2D

⑤ Rototranslation - 3D

⑥ Composition

⑦ Projective 2D Geometry

⑧ Projective Transformations



Projective Transformations - Definition

Definition

A projectivity is an invertible mapping $h(\cdot) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

x_1, x_2, x_3 lie on the same line $\iff h(x_1), h(x_2), h(x_3)$ do

i.e., a projectivity maintains collinearity

Theorem

A mapping $h(\cdot) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a projectivity

1

\exists a non-singular 3×3 matrix \mathbf{H} such that

$\forall \mathbf{p} \in \mathbb{R}^2$ expressed with its homogeneous vector \mathbf{p}_h

$$h(\mathbf{p}_h) = \mathbf{H} \mathbf{p}_h$$

Projective Transformations - Practice

PROJECTIVE TRANSFORMATION

$$\mathbf{x}' = \mathbf{H}\mathbf{x}$$

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Projective Transformations - Practice

PROJECTIVE TRANSFORMATION

$$\mathbf{x}' = \mathbf{H}\mathbf{x}$$

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

NOTE

- \mathbf{H} has 9 elements
- \mathbf{H} is homogeneous too: $\lambda\mathbf{H} \equiv \mathbf{H}$

normalized if $h_{33} = 1$

- → only 8 D.O.F.

Projective Transformations - Practice

PROJECTIVE TRANSFORMATION

$$\mathbf{x}' = \mathbf{H}\mathbf{x}$$

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

NOTE

- \mathbf{H} has 9 elements
- \mathbf{H} is homogeneous too: $\lambda\mathbf{H} \equiv \mathbf{H}$

normalized if $h_{33} = 1$

- \rightarrow only 8 D.O.F.

SYNONYMOUS


- Projectivity
- *Projective transformation*
- Collineation
- *Homography*

Projective Transformations - Mapping between planes

ORIGINAL IMAGE



RECTIFIED IMAGE



ESTIMATION

- Take four point on first image \mathbf{x}_i
- Map on four known destination points \mathbf{x}'_i

- Solve
$$\begin{cases} \mathbf{x}'_1 = \mathbf{H}\mathbf{x}_1 \\ \mathbf{x}'_2 = \mathbf{H}\mathbf{x}_2 \\ \mathbf{x}'_3 = \mathbf{H}\mathbf{x}_3 \\ \mathbf{x}'_4 = \mathbf{H}\mathbf{x}_4 \end{cases} \text{ on } h_{ij}.$$

NOTE

- \mathbf{H} has 8 D.O.F. ($\lambda\mathbf{H} = \mathbf{H}$)
- each point impose 2 constraint

Details

*“Multiple View Geometry
in computer vision”
Hartley Zisserman
Chapter 4.*